

Macroscopic quantum spin tunnelling with two interacting spins.

Solomon A. Owerre* and M. B. Paranjape†

*Groupe de physique des particules, Département de physique, Université de Montréal,
C.P. 6128, succ. centre-ville, Montréal, Québec, Canada, H3C 3J7*

ABSTRACT

We study the simple Hamiltonian, $H = -K(S_{1z}^2 + S_{2z}^2) + \lambda \vec{S}_1 \cdot \vec{S}_2$, of two, large, coupled spins which are taken equal, each of total spin s . The exact ground state of this simple Hamiltonian is not known for an antiferromagnetic coupling, $\lambda > 0$. In the absence of the exchange interaction, the ground state is four fold degenerate, corresponding to the states where the individual spins are in their highest weight or lowest weight states, $|\uparrow, \uparrow\rangle, |\downarrow, \downarrow\rangle, |\uparrow, \downarrow\rangle, |\downarrow, \uparrow\rangle$, in obvious notation. The first two remain exact eigenstates of the full Hamiltonian. However, we show that the two states $|\uparrow, \downarrow\rangle, |\downarrow, \uparrow\rangle$ organize themselves into the combinations $|\pm\rangle = \frac{1}{\sqrt{2}}(|\uparrow, \downarrow\rangle \pm |\downarrow, \uparrow\rangle)$, up to perturbative corrections. For $\lambda > 0$, the ground state is non-degenerate, and we find the interesting result that for integer spins the ground state is $|+\rangle$, and the first excited state is the anti-symmetric combination $|-\rangle$ while for half odd integer spin, these roles are exactly reversed. The energy splitting however, is proportional to λ^{2s} , as expected by perturbation theory to the $2s^{\text{th}}$ order. We obtain these results through the spin coherent state path integral.

PACS numbers: 73.40.Gk, 75.45.+j, 75.50.Ee, 75.50.Gg, 75.50.Xx, 75.75.Jn

Introduction- We study the case of two large, coupled, quantum spins in the presence of a large, simple, easy axis anisotropy, interacting with each other through a standard spin-spin exchange coupling, corresponding to the Hamiltonian

$$H = -K(S_{1z}^2 + S_{2z}^2) + \lambda \vec{S}_1 \cdot \vec{S}_2. \quad (1)$$

We will consider $K > 0$ and specialize to the case of equal spins $\vec{S}_1 = \vec{S}_2 = \vec{S}$. $\lambda > 0$ gives anti-ferromagnetic coupling while $\lambda < 0$ is ferromagnetic. The spins $\vec{S}_i$ could correspond to quantum spins of macroscopic multi-atomic molecules [1–3], or the quantum spins a macroscopic ferromagnetic grains [4], or the average spin of each of the two staggered Néel sub-lattices in a quantum anti-ferromagnet [2, 5].

The non-interacting system is defined by $\lambda = 0$, here the spin eigenstates of S_{iz} , notationally $|s, s_{1z}\rangle \otimes |s, s_{2z}\rangle \equiv |s_{1z}, s_{2z}\rangle$, are obviously exact eigenstates. The ground state is four-fold degenerate, corresponding to the states $|s, s\rangle, |-s, -s\rangle, |s, -s\rangle$ and $|-s, s\rangle$, which we will write as $|\uparrow, \uparrow\rangle, |\downarrow, \downarrow\rangle, |\uparrow, \downarrow\rangle, |\downarrow, \uparrow\rangle$, each with energy $E = -2Ks^2$. The first excited state, which is 8 fold degenerate, is split from the ground state by energy $\Delta E = K(2s - 1)$.

In the weak coupling limit, $\lambda/K \rightarrow 0$, it is an interesting question to ask what is the ground state and the first few excited states of the system for large spin $\vec{S}$. Surprisingly, this is a yet in general unsolved problem. For spin $1/2$, the exact eigenstates are trivially found, for spin 1 , the problem is a 9×9 matrix, which again can be diagonalized, but soon the problem becomes intractable. In principle we must diagonalize a $(2s + 1)^2 \times (2s + 1)^2$ matrix, that though is rather sparse, is not amenable to an exact diagonalization. For weak coupling the anisotropic potential continues to align or anti-align the

spins along the z axis in the ground state.

As the non-interacting ground state is four fold degenerate, in first order degenerate perturbation theory, we should diagonalize the exchange interaction in the degenerate subspace. However, it turns out to be already diagonal in that subspace. The exact eigenvalue is $\pm \lambda s^2$ for the two states $|\uparrow, \uparrow\rangle, |\downarrow, \downarrow\rangle$ and then the full Hamiltonian, Eqn.(1), has total energy eigenvalue $(-2K \pm \lambda)s^2$. The full Hamiltonian can be alternatively written as

$$H = -K(S_{1z}^2 + S_{2z}^2) \pm \lambda \left(S_{1z}S_{2z} + \frac{1}{2}(S_1^+ S_2^- + S_1^- S_2^+) \right), \quad (2)$$

where $S_i^\pm = S_{ix} \pm iS_{iy}$. $S_i^\pm$ act as raising and lowering operators for S_{iz} . Thus they annihilate the states $|\uparrow, \uparrow\rangle, |\downarrow, \downarrow\rangle$ respectively. For the two states $|\uparrow, \downarrow\rangle, |\downarrow, \uparrow\rangle$ the perturbation continues to be diagonal but only in the degenerate subspace, with eigenvalue $\mp \lambda s^2$ yielding, in first order degenerate perturbation theory, a total energy of $(-2K \mp \lambda)s^2$.

However, we will find that the states $|\pm\rangle = \frac{1}{\sqrt{2}}(|\uparrow, \downarrow\rangle \pm |\downarrow, \uparrow\rangle)$ are the appropriate linear combinations implied by degenerate perturbation theory. The perturbing Hamiltonian links the state $|\pm s, \mp s\rangle$ only to the state $|\pm s \mp 1, \mp s \pm 1\rangle$. To reach the state $|\mp s, \pm s\rangle$ from the state $|\pm s, \mp s\rangle$ requires one to go to $2s^{\text{th}}$ order in perturbation, and s is assumed to be large. Indeed, we find our results via macroscopic quantum tunnelling using the spin coherent state path integral. Using the path integral to determine large orders in perturbation theory has already been studied in field theory [6].

Spin coherent state path integral - The quantum (large) spin systems can be described by the spin coherent state

path integral [10, 11].

$$\langle \chi | e^{-\beta H} | \psi \rangle = \mathcal{N} \int_{\psi}^{\chi} \mathcal{D}\theta_i \mathcal{D}\phi_i e^{-S_E}. \quad (3)$$

S_E is the Euclidean action which corresponding to dynamics of particles moving on a two sphere, and which contains first order kinetic term, the Wess-Zumino-Novikov-Witten (WZNW) term for the spin degree of freedom [7].

Our two spin system, in real time, is governed by an action $S = \int dt \mathcal{L}$ where,

$$\begin{aligned} \mathcal{L} = & \int dx \sigma_1 \hat{S}_1 \cdot (\partial_t \hat{S}_1 \times \partial_x \hat{S}_1) - V_1(\hat{S}_1) + \\ & + \int dx \sigma_2 \hat{S}_2 \cdot (\partial_t \hat{S}_2 \times \partial_x \hat{S}_2) - V_2(\hat{S}_2) - \lambda \hat{S}_1 \cdot \hat{S}_2 \end{aligned} \quad (4)$$

where now $\hat{S}_i = (\sin \theta_i \cos \phi_i, \sin \theta_i \sin \phi_i, \cos \theta_i)$, $i = 1, 2$ are three vectors of unit norm, representing semi-classically the quantum spin [4] and σ_i are the values of each spin. In terms of spherical coordinates the Lagrangian takes the form

$$\begin{aligned} \mathcal{L} = & -\sigma_1 \dot{\phi}_1 (1 - \cos \theta_1) - V_1(\theta_1, \phi_1) \\ & - \sigma_2 \dot{\phi}_2 (1 - \cos \theta_2) - V_2(\theta_2, \phi_2) \\ & - \lambda (\sin \theta_1 \sin \theta_2 \cos(\phi_1 - \phi_2) + \cos \theta_1 \cos \theta_2). \end{aligned} \quad (5)$$

The coefficients of the WZNW terms σ_1 and σ_2 are quantized to be a half integer, $\sigma_i = N_i/2$.

We consider the special case of equal spins, with $\sigma_1 = \sigma_2 = s$. Our analysis is valid if we restrict our attention to an external potential with easy-axis, azimuthal symmetry, with a reflection symmetry (along the azimuthal axis), as in [8], $V_i(\theta_i, \phi_i) \equiv V(\theta_i) = V(\pi - \theta_i)$, $i = 1, 2$. The potential is further assumed to have a minimum at the north pole and the south pole, at $\theta_i = 0$, and π . In our case the potential is explicitly

$$V(\hat{S}_i) \equiv V(\theta_i, \phi_i) = K \sin^2 \theta_i. \quad (6)$$

It was shown in Ref. [8], for uncoupled spins, that quantum tunnelling between the spin up and down states of each spin separately is actually absent because of conservation of the z component of each spin. With the exchange interaction only the total z component is conserved allowing transitions $|\uparrow, \downarrow\rangle \longleftrightarrow |\downarrow, \uparrow\rangle$. In general tunnelling exists if there is an equipotential path that links the beginning and end points. We will see that such an equipotential path exists, but through complex values of the phase space variables.

We must find the critical points of the Euclidean action with $t \rightarrow -i\tau$, see Ref. [9], which gives

$$\begin{aligned} \mathcal{L}_E = & is\dot{\phi}_1 (1 - \cos \theta_1) + V(\theta_1) + is\dot{\phi}_2 (1 - \cos \theta_2) + V(\theta_2) \\ & + \lambda (\sin \theta_1 \sin \theta_2 \cos(\phi_1 - \phi_2) + \cos \theta_1 \cos \theta_2). \end{aligned} \quad (7)$$

The solutions must start at $(\theta_1, \phi_1) = (0, 0)$ and $(\theta_2, \phi_2) = (\pi, 0)$, say, and evolve to $(\theta_1, \phi_1) = (\pi, 0)$ and $(\theta_2, \phi_2) = (0, 0)$. In Euclidean time, the WZNW term has become imaginary and the equations of motion in general only have solutions for complexified field configurations. Varying with respect to ϕ_i gives equations that correspond to conservation of angular momentum

$$is \frac{d}{d\tau} (1 - \cos \theta_1) + \lambda \sin \theta_1 \sin \theta_2 \sin(\phi_1 - \phi_2) = 0 \quad (8)$$

$$is \frac{d}{d\tau} (1 - \cos \theta_2) - \lambda \sin \theta_1 \sin \theta_2 \sin(\phi_1 - \phi_2) = 0 \quad (9)$$

While varying with respect to θ_i gives the equations:

$$\frac{\partial \mathcal{L}_E}{\partial \theta_1} = 0 = \frac{\partial \mathcal{L}_E}{\partial \theta_2} \quad (10)$$

Adding Eqn's (8) and (9) we simply get

$$\frac{d}{d\tau} (\cos \theta_1 + \cos \theta_2) = 0. \quad (11)$$

Hence

$$\cos \theta_1 + \cos \theta_2 = l = 0 \implies \theta_2 = \pi - \theta_1 \quad (12)$$

where the constant l is chosen to be zero using the initial condition $\theta_1 = 0, \theta_2 = \pi$. We can now eliminate θ_2 from the equations of motion and writing $\theta = \theta_1$, $\phi = \phi_1 - \phi_2$ and $\Phi = \phi_1 + \phi_2$ and taking $V_i(\theta_i) = V(\theta_i) = V(\pi - \theta_i)$ we get the effective Lagrangian:

$$\mathcal{L} = is\dot{\Phi} - is\dot{\phi} \cos \theta + U(\theta, \phi) \quad (13)$$

where

$$U(\theta, \phi) = 2V(\theta) + \lambda (\sin^2 \theta \cos \phi - \cos^2 \theta) + \lambda \quad (14)$$

is the effective potential energy, where we have removed a constant λ so that the potential is normalized to zero at $\theta = 0$. The first term in the Lagrangian is a total derivative and drops out. The equations of motion become:

$$is\dot{\phi} \sin \theta = - \frac{\partial U(\theta, \phi)}{\partial \theta} \quad (15)$$

$$is\dot{\theta} \sin \theta = \frac{\partial U(\theta, \phi)}{\partial \phi} \quad (16)$$

These equations have no solutions on the space of real functions $\theta(\tau), \phi(\tau)$ due to the explicit i on the left hand side. The analog of conservation of energy follows immediately from these equations, this is easily derived by multiplying (15) by $\dot{\theta}$ and (16) by $\dot{\phi}$ and subtracting, this gives:

$$\frac{dU(\theta, \phi)}{d\tau} = 0 \quad \text{i.e.,} \quad U(\theta, \phi) = \text{const.} = 0 \quad (17)$$

The constant has been set to 0 using the initial condition $\theta = 0$. From Eqn. (14), we have

$$U(\theta, \phi) = (2K + \lambda(\cos \phi + 1)) \sin^2 \theta = 0 \quad (18)$$

implying $(2K + \lambda(\cos \phi + 1)) = 0$ as $\sin^2 \theta \neq 0$, which is required for a non-trivial solution. Then

$$\cos \phi = -\left(\frac{2K}{\lambda} + 1\right) \quad (19)$$

and we see that ϕ must be a constant. This is not valid in general, it is due to the specific choice of the external potential Eqn. (6). Since $K > |\lambda|$ we get $|\cos \phi| > 1$, which of course has no solution for real ϕ . We take $\phi = \phi_R + i\phi_I$ which gives $\cos \phi = \cos \phi_R \cosh \phi_I - i \sin \phi_R \sinh \phi_I$. As the RHS of Eqn. (19) is real, we must have either $\phi_I = 0$ or $\phi_R = n\pi$ or both. Clearly the $\phi_I = 0$ cannot yield a solution for Eqn. (19), hence we must have $\phi_R = n\pi$. As we must impose 2π periodicity on ϕ_R only $n = 0$ or 1 exist. Then we get

$$\cos \phi = (-1)^n \cosh \phi_I = \begin{cases} -\left(\frac{2K}{\lambda} + 1\right) & \text{if } \lambda > 0 \\ +\left(\frac{2K}{|\lambda|} - 1\right) & \text{if } \lambda < 0 \end{cases} \quad (20)$$

Thus $n = 1$ for $\lambda > 0$ and $n = 0$ for $\lambda < 0$ allowing for the unified expression

$$\cosh \phi_I = \frac{2K + \lambda}{|\lambda|}. \quad (21)$$

Eqn. (16) simplifies to

$$is \frac{\dot{\theta}}{\sin \theta} = -\lambda \sin \phi = -i\lambda(-1)^n \sinh \phi_I = i|\lambda| \sinh \phi_I \quad (22)$$

as $\lambda(-1)^n = -|\lambda|$. Eqn. (21) has two solutions positive ϕ_I corresponds to the instanton, ($\dot{\theta} > 0$), and negative ϕ_I corresponds to the anti-instanton, ($\dot{\theta} < 0$). The equation is trivially integrated with solution

$$\theta(\tau) = 2 \arctan \left(e^{\omega(\tau - \tau_0)} \right) \quad (23)$$

where $\omega = (|\lambda|/s) \sinh \phi_I$ and at $\tau = \tau_0$ we have $\theta(\tau) = \pi/2$. Thus $\theta(\tau)$ interpolates from 0 to π as $\tau = -\infty \rightarrow \infty$ for the instanton and from π to 0 for an anti-instanton.

Using $\dot{\phi} = 0$ and (17) we see that the action for this instanton trajectory, let us call it S_0 , simply vanishes

$$S_0 = \int_{-\infty}^{\infty} d\tau \mathcal{L} = 0. \quad (24)$$

So where could the amplitude come from? We have not taken into account the fact that ϕ must be translated from $\phi = 0$ (any initial point will do, as long as it is consistently used to compute the full amplitude) to $\phi = n\pi + i\phi_I$ before the instanton can occur and then back to $\phi = 0$ after the instanton has occurred. Normally such a translation has no effect, either the change

at the beginning cancels that at the end, or if the action is second order in time derivative, moving adiabatically gives no contribution. But in the present case, before the instanton occurs, $\theta = 0$, but after it has occurred, $\theta = \pi$. As $\dot{\phi}$ is multiplied by $\cos \theta$ in the action, the two contributions actually add, there is a net contribution to the action. Indeed the change of the full action for the combination of the instanton and the changes in ϕ is given by

$$\begin{aligned} \Delta S &= \int_0^{n\pi + i\phi_I} -isd\phi \cos \theta|_{\theta=0} + S_0 + \int_{n\pi + i\phi_I}^0 -isd\phi \cos \theta|_{\theta=\pi} \\ &= -is2n\pi + 2s\phi_I \end{aligned} \quad (25)$$

We will use this information to compute the following matrix element, using the spin coherent states $|\theta, \phi\rangle$ and the lowest two energy eigenstates $|E_0\rangle$ and $|E_1\rangle$:

$$\begin{aligned} \langle \theta_f, \phi_f | e^{-\beta H} | \theta_i, \phi_i \rangle &= e^{-\beta E_0} \langle \theta_f, \phi_f | E_0 \rangle \langle E_0 | \theta_i, \phi_i \rangle \\ &+ e^{-\beta E_1} \langle \theta_f, \phi_f | E_1 \rangle \langle E_1 | \theta_i, \phi_i \rangle + \dots \end{aligned} \quad (26)$$

On the other hand, the matrix element is given by the spin coherent state path integral

$$\langle \theta_f, \phi_f | e^{-\beta H} | \theta_i, \phi_i \rangle = \mathcal{N} \int_{\theta_i, \phi_i}^{\theta_f, \phi_f} \mathcal{D}\theta \mathcal{D}\phi e^{-S_E}. \quad (27)$$

The integration is done in the saddle point approximation. With $(\theta_i, \phi_i) = (0, 0)$ corresponding to the state $|\uparrow, \downarrow\rangle$ and $(\theta_f, \phi_f) = (\pi, 0)$ corresponding to the state $|\downarrow, \uparrow\rangle$ we get, with a mild abuse of notation

$$\langle \downarrow, \uparrow | e^{-\beta H} | \uparrow, \downarrow \rangle = \mathcal{N} e^{-\Delta S \kappa \beta} (1 + \dots) \quad (28)$$

where κ is the ratio of the square root of the determinant of the operator governing the second order fluctuations, except for the time translation zero mode, about the instanton and that of the free determinant. It can in principle be calculated, but we have not done this. The zero mode is taken into account by integrating over the position of the occurrence of the instanton giving rise to the factor of β . $\mathcal{N}$ is the overall normalisation including the square root of the free determinant which is given by $N e^{-E_0 \beta}$ where E_0 is the unperturbed ground state energy and N is a constant from the ground state wave function. The result exponentiates, but since we must sum over all sequences of one instanton followed by any number of anti-instanton instanton pairs the total number of instantons and anti-instantons is odd, and we get

$$e^{-\Delta S \kappa \beta} \rightarrow \sinh(e^{-\Delta S \kappa \beta}) \quad (29)$$

Given $\Delta S = -is2n\pi + 2s\phi_I$ and solving Eqn. (21) for ϕ_I for $K \gg |\lambda|$

$$\phi_I = \text{arccosh} \left(\frac{2K + \lambda}{|\lambda|} \right) \approx \ln \left(\frac{4K}{|\lambda|} \right) \quad (30)$$

gives [12]:

$$e^{-\Delta S} = \begin{cases} e^{is2\pi - 2s\phi_I} & \text{if } \lambda > 0 = \begin{cases} \left(\frac{|\lambda|}{4K}\right)^{2s} & \text{if } s \in \mathbf{Z} \\ -\left(\frac{|\lambda|}{4K}\right)^{2s} & \text{if } s \in \mathbf{Z} + 1/2 \end{cases} \\ \left(\frac{|\lambda|}{4K}\right)^{2s} & \text{if } \lambda < 0 \end{cases} \quad (31)$$

Then we get

$$\langle \downarrow, \uparrow | e^{-\beta H} | \uparrow, \downarrow \rangle = \pm \left(\frac{1}{2} e^{\left(\frac{|\lambda|}{4K}\right)^{2s} \kappa \beta} - \frac{1}{2} e^{-\left(\frac{|\lambda|}{4K}\right)^{2s} \kappa \beta} \right) N e^{-\beta E_0} \quad (32)$$

where the $-$ sign only applies for the case of anti-ferromagnetic coupling with half odd integer spin, $\lambda > 0, s = \mathbf{Z} + 1/2$. An essentially identical analysis yields

$$\begin{aligned} \langle \downarrow, \uparrow | e^{-\beta H} | \downarrow, \uparrow \rangle &= \langle \uparrow, \downarrow | e^{-\beta H} | \uparrow, \downarrow \rangle \\ &= \left(\frac{1}{2} e^{\left(\frac{|\lambda|}{4K}\right)^{2s} \kappa \beta} + \frac{1}{2} e^{-\left(\frac{|\lambda|}{4K}\right)^{2s} \kappa \beta} \right) N e^{-\beta E_0}. \end{aligned} \quad (33)$$

These calculated matrix elements should now be compared with what is expected for the exact theory:

$$\begin{aligned} \langle \downarrow, \uparrow | e^{-\beta H} | \uparrow, \downarrow \rangle &= e^{-\beta(E_0 - \frac{1}{2}\Delta E)} \langle \downarrow, \uparrow | E_0 \rangle \langle E_0 | \uparrow, \downarrow \rangle \\ &+ e^{-\beta(E_0 + \frac{1}{2}\Delta E)} \langle \downarrow, \uparrow | E_1 \rangle \langle E_1 | \uparrow, \downarrow \rangle \end{aligned} \quad (34)$$

and say

$$\begin{aligned} \langle \downarrow, \uparrow | e^{-\beta H} | \downarrow, \uparrow \rangle &= e^{-\beta(E_0 - \frac{1}{2}\Delta E)} \langle \downarrow, \uparrow | E_0 \rangle \langle E_0 | \downarrow, \uparrow \rangle \\ &+ e^{-\beta(E_0 + \frac{1}{2}\Delta E)} \langle \downarrow, \uparrow | E_1 \rangle \langle E_1 | \downarrow, \uparrow \rangle \end{aligned} \quad (35)$$

The energy splitting can be read off from this result

$$\Delta E = E_1 - E_0 = 2 \left(\frac{|\lambda|}{4K} \right)^{2s} \kappa \quad (36)$$

and our main result follows, the low energy eigenstates are given by

$$|E_0\rangle = \frac{1}{\sqrt{2}}(|\downarrow, \uparrow\rangle + |\uparrow, \downarrow\rangle) \quad |E_1\rangle = \frac{1}{\sqrt{2}}(|\downarrow, \uparrow\rangle - |\uparrow, \downarrow\rangle) \quad (37)$$

for $\lambda < 0$ (although here the energy eigenstates are should be $|E_3\rangle$ and $|E_4\rangle$) and $\lambda > 0$ for $s \in \mathbf{Z}$, while for the anti-ferromagnetic $\lambda > 0$ case with half odd integer spin we get

$$|E_0\rangle = \frac{1}{\sqrt{2}}(|\downarrow, \uparrow\rangle - |\uparrow, \downarrow\rangle) \quad |E_1\rangle = \frac{1}{\sqrt{2}}(|\downarrow, \uparrow\rangle + |\uparrow, \downarrow\rangle). \quad (38)$$

This understanding of the ground state in the anti-ferromagnetic case is our main result.

Conclusions- We have found the low energy eigenvalues and the corresponding eigenstates for the Hamiltonian of two equal, large, spins interacting with an easy axis anisotropy and a standard exchange interaction, the latter which is considered as a perturbation. We find that the two states $|\downarrow, \uparrow\rangle, |\uparrow, \downarrow\rangle$ reorganize into the symmetric and the anti-symmetric superposition because of quantum tunnelling transitions. These transitions correspond to the $2s^{\text{th}}$ order effects in perturbation theory. The symmetric combination is the lower energy state for integer spin while the anti-symmetric state is the the lower energy state for half odd integer spins. These states are respectively the ground states for an anti-ferromagnetic coupling.

Acknowledgments- We thank Ian Affleck, Sung-Sik Lee and Cliff Burgess for useful discussions and we thank NSERC of Canada for financial support.

* Electronic address: solomon.akaraka.owerre@umontreal.ca

† Electronic address: paranj@lps.umontreal.ca

- [1] J. Ummethum, J. Nehrkorn, S. Mukherjee, N. B. Ivanov1, S. Stuiber, Th. Strässle, P. L. W. Tregenna-Piggott, H. Mutka, G. Christou, O. Waldmann, and J. Schnack Phys. Rev. B **86**, 104403 (2012)
- [2] O. Waldmann, C. Dobe, H. U. Güdel, and H. Mutka Phys. Rev. B **76**, 054429 (2006)
- [3] Florian Meier, Jeremy Levy and Daniel Loss, Phys. Rev. B **68**, 134417 (2003)
- [4] E.M. Chudnovsky and L. Gunther, Phys. Rev. Lett. **60**, 661(1988); Phys. Rev. B **37**, 9455 (1988).
- [5] Florian Meier and Daniel Loss Phys. Rev. Lett. **86**, 5373 (2001); Phys. Rev. B **64**, 224411 (2001)
- [6] J. C. Collins and D. E. Soper, Annals Phys. **112**, 209 (1978); J. Zinn-Justin, Physics Reports **70**, 109 (1981)
- [7] J. Wess, B. Zumino, Phys. Lett. B **37**:95,1971; S.P. Novikov, Usp.Mat.Nauk, 37N5:3-49,1982; E. Witten, Nucl. Phys., B**160**:57,1979.
- [8] Avinash Khare, M.B Paranjape, Phys. Rev. B **83**, 172401 (2011) [arXiv:hep-th/0806.4692].
- [9] Sidney Coleman Phys. Rev. D **15**, 10 (1977); Anupam Garg, Evgueny Kochetov, Kee-Su Park and Michael Stone, Journal of mathematical physics, **44**, 48 (2003)
- [10] D.Loss, D. P. Di Vincenzo, G. Grinstein, Phys. Rev. Lett. **69**, 3232 (1992); J. von Delft, C. Henley, Phys. Rev. Lett. **69**, 3236(1992)
- [11] John R. Klauder Phys. Rev. D **19**, 2349 (1978)
- [12] We thank Sung-Sik Lee for a discussion at this point which led to the understanding of the (expected to him) interesting difference between integer and half odd integer spins for the anti-ferromagnetic case in our calculation.